

AD-A175 060

NON-ORTHOGONAL DESIGNS FOR MEASURING DISPERSION(U)
CALIFORNIA UNIV RIVERSIDE DEPT OF STATISTICS S GHOSH
18 SEP 86 TR-148 AFOSR-TR-86-2130 AFOSR-86-0048

1/1


UNCLASSIFIED

F/G 12/1

NL



END
1-87



NATIONAL BUREAU OF STANDARDS
 RESOLUTION TEST CHART
 1963-A

2

SECURITY

AD-A175 060

DOCUMENTATION PAGE

1a. REPO Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY NA			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; Distribution Unlimited		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE NA					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) Technical Report No. 148			5. MONITORING ORGANIZATION REPORT NUMBER AFOSR-TR- 86-2130		
6a. NAME OF PERFORMING ORGANIZATION University of California Riverside		6b. OFFICE SYMBOL (If applicable) OL	7a. NAME OF MONITORING ORGANIZATION AFOSR/NM		
6c. ADDRESS (City, State and ZIP Code) Department of Statistics University of California, Riverside Riverside, CA 92521			7b. ADDRESS (City, State and ZIP Code) Bldg. 410 Bolling AFB DC 20332-6448		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION AFOSR		8b. OFFICE SYMBOL (If applicable) NM	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER AFOSR-86-0048		
8c. ADDRESS (City, State and ZIP Code) Bldg. 410 Bolling AFB DC 20332-6448			10. SOURCE OF FUNDING NOS.		
			PROGRAM ELEMENT NO. 61102F	PROJECT NO. 2304	TASK NO. A5
11. TITLE (Include Security Classification) Non-orthogonal Designs For Measuring Dispersion					
12. PERSONAL AUTHOR(S) Subir Ghosh					
13a. TYPE OF REPORT Interim		13b. TIME COVERED FROM July 86 TO Sep 86		14. DATE OF REPORT (Yr., Mo., Day) 18 September 1986	
				15. PAGE COUNT 13	
16. SUPPLEMENTARY NOTATION (invited paper in a special volume on factor screening experiments) Submitted to Communications in Statistics					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB. GR.	Dispersion effects, Factor Screening, Linear Models, Search Designs, Sequential Experiments		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) "Dispersion" effects are considered in addition to "Location" effects of factors in the inferential procedure of sequential factor screening experiments with m factors each at two levels under search linear models. Search designs in measuring "Dispersion" and "Location" effects of factors are presented for both stage one and stage two of factor screening experiments with $4 \leq m \leq 10$.					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS <input type="checkbox"/>			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Major Brian W. Woodruff			22b. TELEPHONE NUMBER (Include Area Code) (202)767-5027		22c. OFFICE SYMBOL AFOSR/NM

DTIC
ELECTE
DEC 11 1986
S E D

NON-ORTHOGONAL DESIGNS FOR MEASURING DISPERSION
EFFECTS IN SEQUENTIAL FACTOR SCREENING EXPERIMENTS
USING SEARCH LINEAR MODELS

Subir Ghosh

University of California, Riverside

Accession Number	
NTIS	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution	
Availability Codes	
Avail and/or	
Dist	Special

A1

ABSTRACT

"Dispersion" effects are considered in addition to "Location" effects of factors in the inferential procedure of sequential factor screening experiments with m factors each at two levels under search linear models. Search designs in measuring "Dispersion" and "Location" effects of factors are presented for both stage one and stage two of factor screening experiments with $4 \leq m \leq 10$.



1. INTRODUCTION

In a factor screening experiment the problem is to find a few effective factors from a list of a large number of possible factors influencing the response and screen out all non-effective factors. Watson (1961), Patel (1962) and others considered only "location" effects in their study of influence of factors on response. Taguchi and Wu (1985), Box and Meyer (1986) and others

considered both "location" and "dispersion" effects in the study of influence of factors on response. The concept of "dispersion" effects in factor screening experiments was introduced in Taguchi and Wu (1985) for replicated fractional factorial designs and in Box and Meyers (1986) for unreplicated fractional factorial designs. Srivastava (1975, 1976) introduced the search linear model and showed that factor screening problems can be solved using designs under the search linear model (called search designs) which have smaller numbers of treatments than that in comparable designs in the literature. Ghosh (1979), Ghosh and Avila (1985) presented many such search designs. However, those search designs were constructed to measure the "location" effects only and most of them are unable to measure the "dispersion" effects of factors. In this paper we define the "dispersion" effects of factors under the search linear model and present search designs to measure both the "location" and "dispersion" effects for factor screening experiments.

Throughout this paper, we consider the sequential factor screening experiments with m factors each at two levels under search linear models. At the first stage of the experimentation the problem is to estimate the dispersion effects at two levels of factors under search linear models, to use the dispersion effects in finding the most effective factor out of m factors and finally to determine the optimum level combination of the most effective factor using the "signal to noise" ratio. At the second stage, the problem is to estimate again the dispersion effects at four level combinations of every two factors under search linear models, to find two most effective factors out of m factors in presence of interactions and finally to determine the optimum level combinations of two most effective factors. The process continues until we find all effective factors. In section 2 of this paper we discuss the models and inferential procedures. In

section 3 we present search designs ($4 \leq m \leq 10$) for both stage one and stage two of factor screening experimentation.

2. INFERENCE

In a 2^m factorial experiment, the treatments are denoted by (a_1, \dots, a_m) , $a_i = 0, 1$; the general mean, main effects and two factor interactions are denoted by μ , F_i and $F_i F_j$, respectively; the observation corresponding to the treatment (a_1, \dots, a_m) is denoted by $y(a_1, \dots, a_m)$. The expectation form of the model is

$$E(y(a_1, \dots, a_m)) = \mu + \sum_{i=1}^m b_i F_i + \sum_{i=1}^m \sum_{\substack{j=1 \\ i < j}}^m b_i b_j F_i F_j + \dots, \quad (1)$$

where $b_i = 1$ if $a_i = 1$ and $b_i = -1$ if $a_i = 0$.

Stage 1

At the stage 1 of the experimentation we consider m different models and the i th model is

$$E(y(a_1, \dots, a_i, \dots, a_m)) = \mu + b_i F_i, \quad i=1, \dots, m, \quad (2)$$

$$V(y(a_1, \dots, a_i, \dots, a_m)) = \sigma_i^2(a_i), \quad (3)$$

and the observations are uncorrelated, where the variance $\sigma_i^2(0)$ and $\sigma_i^2(1)$, $i=1, \dots, m$, are unknown constants, called the "dispersion" effects for the factor i . We now consider a fractional factorial design with N_i treatments. (Treatments may or may not be all distinct.) We denote for $u = 0, 1$, \underline{y}_i^u = the vector of N_i^u observations corresponding to the treatments with the i th factor at the level u , \bar{y}_i^u = the simple arithmetic mean of N_i^u observations in \underline{y}_i^u ,

$$s_i^2(u) = \frac{\left(\underline{y}_i^u - \bar{y}_i^u \underline{j}_{N_i^u} \right)' \left(\underline{y}_i^u - \bar{y}_i^u \underline{j}_{N_i^u} \right)}{(N_i^u - 1)},$$

$$s_i^2 = \frac{(N_i^0 - 1) s_i^2(0) + (N_i^1 - 1) s_i^2(1)}{(N_i^0 - 1) + (N_i^1 - 1)}.$$

Note that $N_i = N_i^0 + N_i^1$. We now write (2) and (3) as

$$E \begin{pmatrix} y_i^0 \\ y_i^1 \end{pmatrix} = \begin{pmatrix} \frac{j}{N_i^0} & -\frac{j}{N_i^0} \\ \frac{j}{N_i^1} & \frac{j}{N_i^1} \end{pmatrix} \begin{pmatrix} \mu \\ F_i \end{pmatrix} = \begin{pmatrix} \frac{j}{N_i^0} & 0 \\ 0 & \frac{j}{N_i^1} \end{pmatrix} \begin{pmatrix} \mu - F_i \\ \mu + F_i \end{pmatrix}, \quad (4)$$

$$V \begin{pmatrix} y_i^0 \\ y_i^1 \end{pmatrix} = \begin{pmatrix} \sigma_i^2(0)I_{N_i^0} & 0 \\ 0 & \sigma_i^2(1)I_{N_i^1} \end{pmatrix}, \quad (5)$$

where j is a vector with all elements unity. It can be checked that $E(s_i^2(u)) = \sigma_i^2(u)$ and the generalized least squares (and also the ordinary least squares) estimators of $(\mu - F_i)$ and $(\mu + F_i)$ are $\overline{y_i^0}$ and $\overline{y_i^1}$, respectively. We select the i th factor as the most effective if s_i^2 is a minimum for $i \in \{1, \dots, m\}$. We select the u th level of the factor i as the optimum level if the (square of) "signal to noise" ratio $[\overline{y_i^u} / s_i(u)]^2$ is a maximum.

Stage 2

At the stage 2 of the experimentation we consider $\binom{m}{2}$ different models and the (i, j) th model is

$$E(y(a_1, \dots, a_i, \dots, a_j, \dots, a_m)) = \mu + b_i F_i + b_j F_j + b_i b_j F_i F_j, \quad (6)$$

$$i, j = 1, \dots, m, i \neq j,$$

$$V(y((a_1, \dots, a_i, \dots, a_j, \dots, a_m))) = \sigma_{ij}^2(a_i a_j), \quad (7)$$

and the observations are uncorrelated, where the variances $\sigma_{ij}^2(00)$, $\sigma_{ij}^2(01)$, $\sigma_{ij}^2(10)$ and $\sigma_{ij}^2(11)$ are unknown constants, called the "dispersion" effects for the factors i and j . We now consider a fractional factorial design with N_2 treatments. Note

again that a particular treatment may or may not be replicated in the design. We denote for $u, v=0,1$,

\underline{y}_{ij}^{uv} = the vector of N_{ij}^{uv} observations corresponding to the treatments with the levels of factors i and j are u and v , respectively,

\bar{y}_{ij}^{uv} = the simple arithmetic mean of N_{ij}^{uv} observations in \underline{y}_{ij}^{uv} ,

$$s_{ij}^2(uv) = \frac{\left(\underline{y}_{ij}^{uv} - \bar{y}_{ij}^{uv} \underline{j}_{N_{ij}^{uv}} \right)' \left(\underline{y}_{ij}^{uv} - \bar{y}_{ij}^{uv} \underline{j}_{N_{ij}^{uv}} \right)}{(N_{ij}^{uv} - 1)},$$

$$s_{ij}^2 = \frac{\sum_{u=0}^1 \sum_{v=0}^1 (N_{ij}^{uv} - 1) s_{ij}^2(uv)}{\sum_{u=0}^1 \sum_{v=0}^1 (N_{ij}^{uv} - 1)}$$

Observe that $N_{ij} = N_{ij}^{00} + N_{ij}^{01} + N_{ij}^{10} + N_{ij}^{11}$. We write (6) and (7) as

$$E \begin{pmatrix} \underline{y}_{ij}^{00} \\ \underline{y}_{ij}^{01} \\ \underline{y}_{ij}^{10} \\ \underline{y}_{ij}^{11} \end{pmatrix} = \begin{pmatrix} \underline{j}_{N_{ij}^{00}} & -\underline{j}_{N_{ij}^{00}} & -\underline{j}_{N_{ij}^{00}} & \underline{j}_{N_{ij}^{00}} \\ \underline{j}_{N_{ij}^{01}} & -\underline{j}_{N_{ij}^{01}} & \underline{j}_{N_{ij}^{01}} & -\underline{j}_{N_{ij}^{01}} \\ \underline{j}_{N_{ij}^{10}} & \underline{j}_{N_{ij}^{10}} & -\underline{j}_{N_{ij}^{10}} & -\underline{j}_{N_{ij}^{10}} \\ \underline{j}_{N_{ij}^{11}} & \underline{j}_{N_{ij}^{11}} & \underline{j}_{N_{ij}^{11}} & \underline{j}_{N_{ij}^{11}} \end{pmatrix} \begin{pmatrix} \mu \\ F_i \\ F_j \\ F_i F_j \end{pmatrix}$$

$$= \begin{pmatrix} \underline{j}_{N_{ij}^{00}} & 0 & 0 & 0 \\ 0 & \underline{j}_{N_{ij}^{01}} & 0 & 0 \\ 0 & 0 & \underline{j}_{N_{ij}^{10}} & 0 \\ 0 & 0 & 0 & \underline{j}_{N_{ij}^{11}} \end{pmatrix} \begin{pmatrix} \mu - F_i - F_j + F_i F_j \\ \mu - F_i + F_j - F_i F_j \\ \mu + F_i - F_j - F_i F_j \\ \mu + F_i + F_j + F_i F_j \end{pmatrix} \quad (8)$$

$$V \begin{pmatrix} \underline{y}_{ij}^{00} \\ \underline{y}_{ij}^{01} \\ \underline{y}_{ij}^{10} \\ \underline{y}_{ij}^{11} \end{pmatrix} = \begin{pmatrix} \sigma_{ij}^2(00)I_{N_{ij}^{00}} & 0 & 0 & 0 \\ 0 & \sigma_{ij}^2(01)I_{N_{ij}^{01}} & 0 & 0 \\ 0 & 0 & \sigma_{ij}^2(10)I_{N_{ij}^{10}} & 0 \\ 0 & 0 & 0 & \sigma_{ij}^2(11)I_{N_{ij}^{11}} \end{pmatrix} \quad (9)$$

It can be seen that $E(s_{ij}^2(uv)) = \sigma_{ij}^2(uv)$ and the generalized least squares (and also the ordinary least squares) estimators of $(\mu - F_i - F_j + F_i F_j)$, $(\mu - F_i + F_j - F_i F_j)$, $(\mu + F_i - F_j - F_i F_j)$ and $(\mu + F_i + F_j + F_i F_j)$ are \overline{y}_{ij}^{00} , \overline{y}_{ij}^{01} , \overline{y}_{ij}^{10} and \overline{y}_{ij}^{11} , respectively. We select the factors i and j as the most effective factors if s_{ij}^2 is a minimum for i, j in $\{1, \dots, m\}$. We select the level (u, v) of the factors i and j as the optimum level if the (square of) "signal to noise" ratio $[\overline{y}_{ij}^{uv}/s_{ij}^{(uv)}]^2$ is a maximum.

We stop at the stage 2 if Minimum s_i^2 is very close to Minimum s_{ij}^2 ; otherwise we go to the stage 3. The stage 3 inferential procedure is similar to those in stages 1 and 2.

3. DESIGN

The orthogonal fractional factorial designs are indeed efficient but have restriction on the number of treatments in the design. The designs used in Taguchi and Wu (1985), Box and Meyer (1986) are all classical orthogonal designs. In this section we construct designs to measure the "dispersion" effects defined in section 2 for the stage 1 and stage 2 of the experiment, relaxing the restriction on the number of treatments. We want to have enough observations (at least two!) in measuring the "dispersion" effects, the number of treatments to be small and furthermore the designs to be near orthogonal.

3.1 DESIGNS FOR THE STAGE 1 EXPERIMENT

The design condition obtained in Srivastava (1975) requires that for all i and j , $i \neq j$, i, j in $\{1, \dots, m\}$, μ , F_i and F_j are unbiasedly estimable (u.e.) under the model (1) assuming all other parameters to be zero. We denote a design by a treatment matrix $T(N_1 \times m)$ with rows as treatments and columns as factors.

Theorem 1. A necessary and sufficient condition that a treatment matrix $T(N_1 \times m)$ satisfies the design condition is that for every submatrix $T_1(N_1 \times 2)$ of T , at least three pairs out of the four pairs (00), (01), (10) and (11) appear as rows in T_1 .

Proof. To estimate the three parameters μ , F_i and F_j , we need three independent equations in parameters from (1) under the assumption that all other parameters are exactly equal to zero.

For the submatrix $T_1(N_1 \times 2)$ of T corresponding to the factors i and j , any three rows out of four possible distinct rows (00), (01), (10) and (11) will give three independent equations in μ , F_i and F_j under (1). This completes the proof.

The characterization in Theorem 1 is so simple that the checking can even be done by eye inspection. A result similar to Theorem 1 is also available in Srivastava (1975).

We now present some designs for $4 \leq m \leq 10$ satisfying the condition in Theorem 1. In the Table I, for brevity, we indicate a treatment (i.e., a row) in T by the positions where the level 1 is occurring. The treatment (0, ..., 0) will be denoted by θ . To illustrate this for $m=4$, $T(5 \times 4)$ matrix is given below.

$$T = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 1 & 0 & 1 & 0 \end{bmatrix} .$$

We denote this T in our representation as $(1234, \emptyset, 12, 34, 13)$. Notice that in every column of T , $u(=0,1)$ is appearing either twice or thrice.

TABLE I.
DESIGNS ($4 \leq m \leq 10$) FOR THE STAGE 1 EXPERIMENT

m	Designs
4	1234, \emptyset , 12, 34, 13
5	12345, 12, 13, 24, 35
6	3456, 1256, 1234, 146, 235
7	127, 3456, 1234, 567, 1357, 1234567
8	1278, 3456, 3478, 1256, 1357, 1234
9	1278, 3456, 34789, 1256, 13579, 1234
10	12789 $\overline{10}$, 3456, 34789 $\overline{10}$, 1256, 13579, 1234, 24689

3.2 DESIGNS FOR THE STAGE 2 EXPERIMENT

The design condition obtained in Srivastava (1975) requires for all i, j ($i \neq j$), u, v ($u \neq v$), $(i, j) \neq (u, v)$, i, j, u, v in $\{1, \dots, m\}$, $\mu, F_i, F_j, F_i F_j, F_u, F_v$ and $F_u F_v$ are u.e. under the model (1) assuming all other parameters to be zero.

Theorem 2. Suppose a design $T(N_2 \times m)$ is such that μ, F_i, F_j, F_u are u.e. for every distinct i, j and u in $\{1, \dots, m\}$ under (1) assuming all other parameters except $F_i F_j$ and $F_i F_u$ to be zero. Then $F_i F_j$ and $F_i F_u$ are also u.e.

Proof. It can be seen using Theorem 1 that $F_i F_j + F_i F_u$ and $-F_i F_j + F_i F_u$ are u.e. This completes the proof.

Theorem 3. Suppose a design $T(N_2 \times m)$ is such that μ, F_i, F_j, F_u and F_v are u.e. for every distinct i, j, u and v in $\{1, \dots, m\}$

under (1) assuming all other parameters except $F_i F_j$ and $F_u F_v$ to be zero. Then $F_i F_j$ and $F_u F_v$ are u.e. if and only if for $T_1 (N_2 \times 2)$ corresponding to factors (i,j) and $T_2 (N_2 \times 2)$ corresponding to factors (u,v) there are two distinct rows in T with the number of 1's in T_1 and T_2 as follows.

Row	The number of 1's in	
	T_1	T_2
1 a or b	even	even
	odd	odd
2 a or b	odd	even
	even	odd

Proof. It can be seen that from the rows 1 and 2 that $F_i F_j + F_u F_v$ and $-F_i F_j + F_u F_v$ are u.e. This completes the proof.

3.2.1 CONSTRUCTION OF DESIGNS $m=4$

The treatment matrix consists of 9 treatments as rows; the first eight treatments are solutions of $x_1 + x_2 + x_3 + x_4 = 0$ over the finite field GF(2) and the last treatment is (1000). The treatment matrix is represented as (1234, 0, 12, 34, 13, 24, 14, 23, 1).

The first eight treatments form an orthogonal resolution IV plan. It follows from Theorem 2 that $\mu, F_i, F_j, F_i F_j, F_u, F_i F_u$ are u.e. for any three distinct i, j and u in $\{1, \dots, m\}$. It can be checked that $\mu, F_i, F_j, F_u, F_v, F_i F_j + F_u F_v$ are u.e. for any four distinct integers i, j, u and v. It can be seen that for all rows in the treatment matrix the numbers of 1's in submatrices corresponding to (i,j) and (u,v) are either (even, even) or (odd, odd) and therefore the treatment matrix does not satisfy the conditions in Theorem 3. When the treatment (1000) is added to the treatment matrix, it follows from Theorem 3 that $\mu, F_i, F_j, F_i F_j, F_u, F_v$ and $F_u F_v$ are u.e.

m=5

The treatment matrix consists of 11 treatments as all (1x5) vectors with two elements are unity and the other elements are zero and furthermore the treatment with the levels of all factors are unity.

This is a balanced array of full strength. There is a complete 2^3 factorial experiment w.r.t. any three factors and there is a resolution V plan for a 2^4 factorial experiment w.r.t. any four factors. Thus the design satisfies the design condition.

m=6

Design 1: The treatment matrix consists of 16 treatments as follows:

The 14 treatments are solutions of $x_1+x_2+x_3+x_4=0$ and $x_3+x_4+x_5+x_6=0$ over the finite field GF(2) excluding the treatments (000000) and (111111); and the other two treatments are (10 00 00) and (00 00 01).

It can be seen that all 2^3 distinct treatments are present in rows of T w.r.t. any three factors F_i , F_j and F_u . It thus follows μ , F_i , F_j , $F_i F_j$, F_u and $F_i F_u$ are u.e. for every three distinct factors i , j and u . For four factors of the type (F_1, F_2, F_3, F_4) , (F_1, F_2, F_5, F_6) and (F_3, F_4, F_5, F_6) , we get basically the design for $m=4$ with replications. We get a resolution V plan for a 2^4 factorial experiment w.r.t. any four factors other than (F_1, F_2, F_3, F_4) , (F_1, F_2, F_5, F_6) and (F_3, F_4, F_5, F_6) . Therefore, μ , F_i , F_j , $F_i F_j$, F_u , F_v and $F_u F_v$ are u.e. for every four distinct factors i , j , u and v .

Design 2: The treatment matrix consists of 15 treatments as follows.

The 15 treatments are solutions of $x_1+x_2+x_3+x_4=0$ and $x_3+x_4+x_5+x_6=0$ over the finite field GF(2) excluding the treatments (01 01 01), (10 10 10) and (00 00 00); and the other two treatments are (10 00 00) and (00 00 01).

The argument is similar to that in Design 1.

m=7

The treatment matrix consists of 17 treatments as follows.

The 14 treatments are solutions of $x_1+x_2+x_3+x_4=0$, $x_3+x_4+x_5+x_6=0$ and $x_1+x_3+x_5+x_7=0$ over the finite field GF(2) excluding the treatments (1111111) and (0000000); and the other 3 treatments are (10 00 00 0), (00 00 01 0), (00 00 00 1). [One may keep (1111111) in the design although the design is all right without it.]

m=8

The treatment matrix consists of 18 treatments as follows.

The 14 treatments are solutions of $x_1+x_2+x_3+x_4=0$, $x_3+x_4+x_5+x_6=0$, $x_5+x_6+x_7+x_8=0$ and $x_1+x_3+x_5+x_7=0$ over the finite field GF(2) excluding the treatments (00 00 00 00) and (11 11 11 11); and the other 4 treatments are (10 00 00 00), (00 00 10 00), (00 00 00 10), (00 00 00 01).

m=9

The treatment matrix consists of 28 treatments as follows.

The treatments are solutions of $x_1+x_2+x_3+x_4=0$, $x_3+x_4+x_5+x_6=0$, $x_5+x_6+x_7+x_8=0$ and $x_1+x_3+x_5+x_7=0$ over the finite field GF(2) excluding the treatments (000000000), (000000001), (11 11 11 11 1), (11 11 11 11 0).

m=10

The treatment matrix consists of 35 treatments as follows.

The 30 treatments are solutions of $x_1+x_2+x_3+x_4=0$, $x_3+x_4+x_5+x_6=0$, $x_5+x_6+x_7+x_8=0$, $x_7+x_8+x_9+x_{10}=0$ and $x_1+x_3+x_5+x_7=0$ over the finite field GF(2) excluding the treatments (00 00 00 00 00), (11 11 11 11 11), and the other five treatments are (10 00 00 00 00), (00 00 10 00 00), (00 00 00 10 00), (00 00 00 00 10), (00 00 00 01 00).

The arguments for the cases $m=7, 8, 9, 10$ are similar to the argument for $m=6$.

4. FINAL REMARKS

This paper touches the major developments in factor screening designs over the period of more than 25 years and deals with the important issue of measuring the "dispersion" effects in addition to the "location" effects of factors using search linear models. The influence of Professor R. C. Bose to this author is not easy to describe in words and this author takes pride in dedicating this work to honor Professor R. C. Bose.

ACKNOWLEDGEMENTS

The work of the author is sponsored by the Air Force Office of Scientific Research under Grant AFOSR-86-0048.

BIBLIOGRAPHY

- Bose, R. C. (1947). Mathematical theory of symmetrical factorial designs. Sankhyā, 8, 107-166.
- Box, G. E. P. and Meyer, R. D. (1986). Dispersion effects from fractional designs. Technometrics, 28, 1927.
- Ghosh, S. (1979). On single and multistage factor screening procedures. Journal of Combinatorics, Information and System Sciences, 4, 275-284.
- Ghosh, S. and Avila, D. (1985). Some new factor screening designs using the search linear model. Journal of Statistical Planning and Inference, 11, 259-266.
- Katona, G. and Srivastava, J. (1983). Minimal 2-coverings of a finite affine space based on GF(2). Journal of Statistical Planning and Inference, 8, 375-388.
- Patel, M. S. (1962). Group-screening with more than two stages. Technometrics, 4, 209-217.
- Srivastava, J. N. (1975). Designs for searching non-negligible effects. In: J. N. Srivastava, Ed., A Survey of Statistical Designs and Linear Models. North-Holland, Amsterdam, 507-519.

- Srivastava, J. N. (1976). Smaller sized factor screening designs through the use of search linear models. 9th Int. Biometric Conference, Boston, August 22-27.
- Taguchi, G. and Wu, Y. (1985). Introduction to off-line quality control. Central Japan Quality Control Association, Tokyo.
- Watson, G. S. (1961). A study of the group screening method. Technometrics, 3, 371-388.

Key Words and Phrases: Dispersion effects, Factor Screening,
Linear models, Search designs, Sequential
experiments.

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)
NOTICE OF TRANSMITTAL TO DTIC
This technical report has been reviewed and is
approved for public release IAW AFR 190-12.
Distribution is unlimited.
MATTHEW J. KERPER

Chief, Technical Information Division

Approved for public release;
distribution unlimited.